

AN OPERATOR FORMULA FOR THE NUMBER OF HALVED MONOTONE TRIANGLES WITH PRESCRIBED BOTTOM ROW

Ilse Fischer

Fakultät für Mathematik, Universität Wien
Nordbergstrasse 15, A-1090 Wien, Austria
E-mail: Ilse.Fischer@univie.ac.at

ABSTRACT. Monotone triangles are certain triangular arrays of integers, which correspond to $n \times n$ alternating sign matrices when prescribing $(1, 2, \dots, n)$ as bottom row of the monotone triangle. In this article we define halved monotone triangles, a specialization of which correspond to vertically symmetric alternating sign matrices. We derive an operator formula for the number of halved monotone triangles with prescribed bottom row which is analogous to our operator formula for the number of ordinary monotone triangles [2].

1. INTRODUCTION

Alternating sign matrices and equivalent objects such as fully packed loop configurations, the six vertex model and monotone triangles are nowadays a rich source for intriguing problems on which combinatorialists can test their various enumeration methods. This article is another contribution in this respect.

In [2] we gave a formula for the number of monotone triangles with prescribed bottom row. Strikingly this formula involves shift operators which are applied to a simple multivariate polynomial. It is an example of a new type of enumeration formula combinatorialists can possibly make use of when answering their enumeration problems. Subsequently, our formula enabled us to give a new proof of the refined alternating sign matrix theorem [3], which was first proved by Zeilberger [12]. Here, we present a second example of such an operator formula. This new formula gives the number of halved monotone triangles with prescribed bottom row, a notion to be defined below.

To keep the treatment self-contained, we recall the basic definitions. An alternating sign matrix is a square matrix with 0s, 1s and -1 s as entries such that the row- and columnsums are 1 and the non-zero entries of each row and of each column alternate in sign. Thus,

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 1 & -1 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

is an alternating sign matrix. The fascinating story of alternating sign matrices [1] began when combinatorialists were confronted with a conjecture by Mills, Robbins and Rumsey [6, 7], which states that the number of $n \times n$ alternating sign matrices is given by the following simple formula

$$\prod_{j=0}^{n-1} \frac{(3j+1)!}{(n+j)!}.$$

For a long time no one could explain this, until finally Zeilberger [11] came up with the first proof. Soon after another, shorter, proof was given by Kuperberg [4]. See also [3], where we have recently presented a new proof of this result.

A *monotone triangle* is a triangular array $(a_{i,j})_{1 \leq j \leq i \leq n}$ of integers,

$$\begin{array}{cccccccc} & & & & & & & a_{1,1} \\ & & & & & & a_{2,1} & a_{2,2} \\ & & & & & a_{3,1} & a_{3,2} & a_{3,3} \\ & & & a_{4,1} & & a_{4,2} & a_{4,3} & a_{4,4} \\ & & a_{5,1} & & a_{5,2} & a_{5,3} & a_{5,4} & a_{5,5} \\ & a_{6,1} & & a_{6,2} & a_{6,3} & a_{6,4} & a_{6,5} & a_{6,6} \\ a_{7,1} & & a_{7,2} & & a_{7,3} & a_{7,4} & a_{7,5} & a_{7,6} & a_{7,7} \end{array}$$

such that $a_{i,j} \leq a_{i-1,j} \leq a_{i,j+1}$ and $a_{i,j} < a_{i,j+1}$ for all i, j . For instance,

$$\begin{array}{cccccccc} & & & & & & & 4 \\ & & & & & & 2 & 6 \\ & & & & 1 & 4 & 7 \\ & & 1 & 3 & 5 & 7 \\ & 1 & 2 & 4 & 6 & 7 \\ 1 & 2 & 3 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{array}$$

is a monotone triangle. It corresponds to the alternating sign matrix above: in the matrix, replace every entry with the sum of entries in the same column above, the entry itself included. The result is a 0-1-matrix with one 1 in the first row, two 1s in the second row etc. If one records the columns of the 1s rowwise, one obtains the corresponding monotone triangle. It is not hard to see that this establishes a bijection between monotone triangles with bottom row $(1, 2, \dots, n)$ and $n \times n$ alternating sign matrices.

Observe that the alternating sign matrix given above is symmetric with respect to the vertical symmetry axis. This is not the case for all alternating sign matrices. In fact there only exist vertically symmetric alternating sign matrices of odd size. (This follows from the fact that an alternating sign matrix has always a unique 1 in its top row.) Kuperberg [5] showed that the number of vertically symmetric $(2n+1) \times (2n+1)$ alternating sign matrices is given by

$$\frac{n!}{(2n)!2^n} \prod_{j=1}^n \frac{(6j-2)!}{(2n+2j-1)!}.$$

(This formula was conjectured by Robbins [10].) The symmetry of vertically symmetric alternating sign matrices translates into a symmetry of the corresponding monotone triangle: the replacement of every entries e by $2n + 2 - e$ and the subsequent reflection along the vertical symmetry axis leaves the monotone triangle invariant. Thus, in case of a vertically symmetric alternating sign matrix, it suffices to “store” (a bit less than) half of the monotone triangle. In our example, this is the following array.

$$\begin{array}{cccc}
 & & & 2 \\
 & & & 1 \\
 & & 1 & 3 \\
 & 1 & 2 & \\
 1 & 2 & 3 &
 \end{array}$$

Note that the middle column of a monotone triangle corresponding to a $(2n + 1) \times (2n + 1)$ vertically symmetric alternating sign matrix consists solely of $(n + 1)$ s and, consequently, we do not have to store it.

These considerations led us to the following definition. A *halved monotone triangle* is a triangular array $(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq \lceil \frac{i}{2} \rceil}$ of integers,

$$\begin{array}{cccc}
 & & & a_{1,1} \\
 & & & a_{2,1} \\
 & & a_{3,1} & a_{3,2} \\
 & a_{4,1} & & a_{4,2} \\
 a_{5,1} & & a_{5,2} & a_{5,3} \\
 a_{6,1} & a_{6,2} & & a_{6,3}
 \end{array}$$

which is monotone increasing in northeast and southeast direction and strictly increasing along rows, that is $a_{i+1,j} \leq a_{i,j}$, $a_{i,j} \leq a_{i+1,j+1}$ and $a_{i,j} < a_{i,j+1}$ for all i, j . The bijection sketched above shows that halved monotone triangles $(a_{i,j})_{1 \leq i \leq 2n, 1 \leq j \leq \lceil \frac{i}{2} \rceil}$ with bottom row $(1, 2, \dots, n)$ such that no entry is greater than n correspond to $(2n+1) \times (2n+1)$ vertically symmetric alternating sign matrices. We are ready to state the main result of the paper.

Theorem 1. *The number of halved monotone triangles with n rows, where no entry exceeds x and with bottom row $(k_1, \dots, k_{\lceil n/2 \rceil})$, $k_1 < k_2 < \dots < k_{\lceil n/2 \rceil} \leq x$, is equal to*

$$\left(\prod_{1 \leq p < q \leq (n+1)/2} E_{k_p} (E_{k_p}^{-1} + E_{k_q}^{-1} - \text{id}) (E_{k_p}^{-1} + E_{k_q} - \text{id}) \right) \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 1 - k_i - k_j)}{(j - i)(j + i - 1)}$$

if n is odd and equal to

$$\left(\prod_{1 \leq p < q \leq n/2} E_{k_p}(E_{k_p}^{-1} + E_{k_q}^{-1} - \text{id})(E_{k_p}^{-1} + E_{k_q} - \text{id}) \right) \prod_{1 \leq i < j \leq n/2} \frac{(k_j - k_i)(2x + 2 - k_i - k_j)}{(j - i)(j + i)} \prod_{i=1}^{n/2} \frac{x + 1 - k_i}{i}$$

if n is even, where $E_x p(x) = p(x + 1)$ denotes the shift operator. In this formula, the product of operators is the composition, and, since the shift operators with respect to different variables commute, we do not have to specify the order in which they are applied.

We think that the following phenomenon is interesting, since it is in analogy to the situation for ordinary monotone triangles, see [2]. If we consider “halved monotone triangles” which do not necessarily have strict monotony along rows (the weak monotony follows from the other conditions), the enumeration problem is significantly easier. These objects are equivalent to shifted plane partitions of trapezoidal shape with prescribed diagonal and were enumerated by Proctor [8, Prop. 4.1]. The number of these halved triangles with n rows, where no entry exceeds x and with bottom row $(k_1, \dots, k_{\lceil n/2 \rceil})$ is equal to

$$\prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i + j - i)(2x + 2 + n - i - j - k_i - k_j)}{(j - i)(j + i - 1)}$$

if n is odd and equal to

$$\prod_{1 \leq i < j \leq n/2} \frac{(k_j - k_i + j - i)(2x + 2 - i - j + n - k_i - k_j)}{(j - i)(j + i)} \prod_{i=1}^{n/2} \frac{x + 1 - i + n/2 - k_i}{i}$$

if n is even. Let $\beta(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ denote the number of these objects. Then, by Theorem 1, the number of halved monotone triangles with n rows, where no entry exceeds x and with bottom row $(k_1, \dots, k_{\lceil n/2 \rceil})$ is given by

$$\left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (E_{k_p} + E_{k_q} - E_{k_p} E_{k_q})(E_{k_p} + E_{k_q}^{-1} - E_{k_p} E_{k_q}^{-1}) \right) \beta(n, x; k_1, \dots, k_{\lceil n/2 \rceil}).$$

This happens to be in perfect analogy to the situation for ordinary monotone triangles: an enumeration formula for the objects with strict monotony along rows can be obtained by applying a product of simple operators to an enumeration formula for the corresponding objects with weak monotony along rows, the latter of which is a simple product formula.

The paper is organized as follows. We prove Theorem 1 in Sections 2 – 5. Our strategy is to first show the polynomiality of the formula, then compute its degree, and finally derive enough properties that characterize the polynomial. To be more precise, in Section 2, we introduce the recursion underlying our enumeration formula for halved

monotone triangles and show the polynomiality of the formula. In Section 3, we define an operator, which is closely related to the recursion and prove a number of lemmas on it. In Section 4, we list and derive the properties that characterize the enumeration polynomial, and, in Section 5, we finally use these properties to prove Theorem 1. In Section 6, we use our operator formulas to derive a generating function for halved monotone triangles as well as a generating function for ordinary monotone triangles.

2. A RECURSION AND THE POLYNOMIALITY OF THE ENUMERATION FORMULA

For $n \geq 1$ and $k_1 < k_2 < \dots < k_{\lceil \frac{n}{2} \rceil} \leq x$, let $\gamma(n, x; k_1, \dots, k_{\lceil \frac{n}{2} \rceil})$ denote the quantity we want to compute, i.e. the number of halved monotone triangles with n rows, where the bottom row is $(k_1, k_2, \dots, k_{\lceil \frac{n}{2} \rceil})$ and all entries are no greater than x . We define a summation operator for functions $f(l_1, \dots, l_{m-1})$, where $m \geq 2$ and $(l_1, \dots, l_{m-1}) \in \mathbb{Z}^{m-1}$, as follows. For given $(k_1, \dots, k_m) \in \mathbb{Z}^m$ we have

$$\sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} := \sum_{\substack{(l_1, \dots, l_{m-1}) \in \mathbb{Z}^{m-1}, \\ k_1 \leq l_1 \leq k_2 \leq \dots \leq k_{m-1} \leq l_{m-1} \leq k_m, l_i \neq l_{i+1}}}, \quad (2.1)$$

i.e. we sum over all strictly increasing sequences $(l_1, \dots, l_{m-1})$ such that $k_i \leq l_i \leq k_{i+1}$ for all i . This operator is well-defined for all strictly increasing sequences $(k_1, \dots, k_m) \in \mathbb{Z}^m$. If we define $\gamma(0, x; -) = 1$ we have the following recursions. If n is even then

$$\gamma(n, x; k_1, \dots, k_{n/2}) = \sum_{(l_1, \dots, l_{n/2})}^{(k_1, \dots, k_{n/2}, x)} \gamma(n-1, x; l_1, l_2, \dots, l_{n/2})$$

and if n is odd then

$$\gamma(n, x; k_1, \dots, k_{(n+1)/2}) = \sum_{(l_1, \dots, l_{(n-1)/2})}^{(k_1, \dots, k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}).$$

We want to extend the interpretation of $\gamma(n, x; k_1, \dots, k_{\lceil \frac{n}{2} \rceil})$ to arbitrary $(k_1, \dots, k_{\lceil \frac{n}{2} \rceil}) \in \mathbb{Z}^{\lceil \frac{n}{2} \rceil}$. For this purpose, it suffices to extend the definition of (2.1) to arbitrary $(k_1, \dots, k_m) \in \mathbb{Z}^m$ and then use the recursions to define the generalization of γ . We use induction with respect to m . For $m = 2$, let

$$\sum_{(l_1)}^{(k_1, k_2)} a(l_1) := \sum_{l_1=k_1}^{k_2} a(l_1),$$

where here and in the following $\sum_{i=a}^b f(i) = - \sum_{i=b+1}^{a-1} f(i)$ if $a > b$. (Note that this implies

$\sum_{i=a}^{a-1} f(i) = 0$. Moreover, $\sum_{x=0}^y p(x)$ will be a polynomial in y if $p(x)$ is a polynomial in x .)

If $m > 2$ we define

$$\begin{aligned} & \sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} a(l_1, \dots, l_{m-1}) := \\ & \sum_{(l_1, \dots, l_{m-2})}^{(k_1, \dots, k_{m-1})} \sum_{l_{m-1}=k_{m-1}+1}^{k_m} a(l_1, \dots, l_{m-2}, l_{m-1}) + \sum_{(l_1, \dots, l_{m-2})}^{(k_1, \dots, k_{m-2}, k_{m-1}-1)} a(l_1, \dots, l_{m-2}, k_{m-1}). \end{aligned}$$

Now it is also obvious that $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is a polynomial in $(k_1, \dots, k_{\lceil n/2 \rceil}, x)$ for fixed n .

This recursion can be used to compute $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ for small values of n . For $n = 1, 2, 3, 4, 5$ you find the results below.

$$\begin{aligned} & 1, x - k_1 + 1, \frac{1}{2}(2x + 2 - k_1 - k_2)(k_2 - k_1 + 1), \frac{1}{6}(x - k_2 + 1)(-k_1^3 + 3xk_1^2 + 6k_1^2 - 2x^2k_1 \\ & + k_2^2k_1 - 10xk_1 - 2xk_2k_1 - 2k_2k_1 - 11k_1 + 2x^2 - xk_2^2 - 2k_2^2 + 7x + 2x^2k_2 + 6xk_2 + 4k_2), \\ & \frac{1}{48}(k_2^2k_1^4 - k_3^2k_1^4 + 2xk_1^4 - 2xk_2k_1^4 - 3k_2k_1^4 + 2xk_3k_1^4 + k_3k_1^4 + 2k_1^4 - 8x^2k_1^3 - 4xk_2^2k_1^3 \\ & - 10k_2^2k_1^3 + 4xk_3^2k_1^3 + 10k_3^2k_1^3 - 28xk_1^3 + 8x^2k_2k_1^3 + 32xk_2k_1^3 + 30k_2k_1^3 - 8x^2k_3k_1^3 - 24xk_3k_1^3 \\ & - 10k_3k_1^3 - 20k_1^3 - k_2^4k_1^2 + k_3^4k_1^2 + 8x^3k_1^2 + 4xk_2^3k_1^2 + 6k_2^3k_1^2 - 4xk_3^3k_1^2 - 2k_3^3k_1^2 + 60x^2k_1^2 + 12xk_2^2k_1^2 \\ & + 24k_2^2k_1^2 - 24xk_3^2k_1^2 - 36k_3^2k_1^2 + 122xk_1^2 - 8x^3k_2k_1^2 - 60x^2k_2k_1^2 - 138xk_2k_1^2 - 99k_2k_1^2 + 8x^3k_3k_1^2 \\ & + 60x^2k_3k_1^2 + 102xk_3k_1^2 + 37k_3k_1^2 + 70k_1^2 + 2xk_2^4k_1 + 5k_2^4k_1 - 2xk_3^4k_1 - 5k_3^4k_1 - 24x^3k_1 - 8x^2k_2^3k_1 \\ & - 32xk_2^3k_1 - 30k_2^3k_1 + 8x^2k_3^3k_1 + 24xk_3^3k_1 + 10k_3^3k_1 - 124x^2k_1 + 8x^3k_2^2k_1 + 36x^2k_2^2k_1 + 42xk_2^2k_1 \\ & + 29k_2^2k_1 - 8x^3k_2^2k_1 - 12x^2k_2^2k_1 + 42xk_2^2k_1 + 31k_2^2k_1 - 152xk_1 + 16x^3k_2k_1 + 96x^2k_2k_1 + 140xk_2k_1 \\ & + 48k_2k_1 - 32x^3k_3k_1 - 144x^2k_3k_1 - 136xk_3k_1 - 36k_3k_1 - 52k_1 - 4xk_2^4 - 6k_2^4 - k_2^2k_3^4 + 2xk_3^4 \\ & + 2xk_2k_3^4 + 3k_2k_3^4 + 4k_3^4 + 16x^2k_2^3 + 48xk_2^3 + 28k_2^3 - 8x^2k_3^3 + 4xk_2^2k_3^3 + 2k_2^2k_3^3 - 20xk_3^3 - 8x^2k_2k_3^3 \\ & - 16xk_2k_3^3 - 6k_2k_3^3 - 8k_3^3 - 16x^3k_2^2 - 72x^2k_2^2 - 100xk_2^2 - 66k_2^2 + k_2^4k_3^2 + 8x^3k_2^2 + 4xk_2^3k_3^2 - 6k_2^3k_3^2 \\ & + 12x^2k_3^2 + 12xk_2^2k_3^2 + 12k_2^2k_3^2 - 22xk_2^2 + 8x^3k_2k_3^2 + 12x^2k_2k_3^2 - 6xk_2k_3^2 + 15k_2k_3^2 - 4k_3^2 + 32x^2k_2 \\ & + 96xk_2 + 44k_2 - 2xk_2^4k_3 - k_2^4k_3 + 24x^3k_3 + 8x^2k_2^3k_3 + 16xk_2^3k_3 + 6k_2^3k_3 + 92x^2k_3 - 8x^3k_2^2k_3 \\ & - 36x^2k_2^2k_3 - 42xk_2^2k_3 - 13k_2^2k_3 + 56xk_3 + 16x^3k_2k_3 + 48x^2k_2k_3 - 4xk_2k_3 - 12k_2k_3 + 8k_3) \end{aligned}$$

This data suggests that the degree of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ in k_i is always $n - 1$. That this is indeed the case will be shown in the following section. However, this comes by surprise because of the following: suppose that $a(l_1, \dots, l_{m-1})$ is a polynomial of degree R in every l_i . Then the degree of

$$\sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} a(l_1, \dots, l_{m-1})$$

in k_i could be as high as $2R + 2$ (e.g. $a(l_1, \dots, l_{m-1}) = \prod_{i=1}^{m-1} l_i^R$). This estimation provides (by induction with respect to n) a bound of $2^{n+1} - 2$ for the degree of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ in k_i .

3. AN OPERATOR RELATED TO THE RECURSION

Most of the definitions and lemmas in this section are taken from [2]. The *shift operator* E_x is defined as $E_x p(x) = p(x + 1)$ and the *difference operator* Δ_x is defined as $E_x - \text{id}$. The *swapping operator* $S_{x,y}$ is defined as $S_{x,y} f(x, y) = f(y, x)$.

Note that the shift operator E_x is invertible as an operator over $\mathbb{C}[x]$, whereas the difference operator Δ_x is not, since it decreases the degree of a polynomial. In the following, we will consider rational functions in shift operators and thus we need a lemma in order to show that the inverses of our denominators exist. (The lemma is a generalization of [2, Lemma 1].) For the statement of the lemma we need the following observation. Let

$$p(X_1, \dots, X_n) = \sum_{(i_1, \dots, i_n) \in (\mathbb{Z}^{\geq 0})^n} a_{i_1, \dots, i_n} X_1^{i_1} \cdots X_n^{i_n}$$

be a formal power series in $(X_1, \dots, X_n)$ over $\mathbb{C}$ and $G(k_1, \dots, k_n)$ be a polynomial in $(k_1, \dots, k_n)$ over $\mathbb{C}$. We define

$$p(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n) := \sum_{(i_1, \dots, i_n) \in (\mathbb{Z}^{\geq 0})^n} a_{i_1, \dots, i_n} \Delta_{k_1}^{i_1} \cdots \Delta_{k_n}^{i_n} G(k_1, \dots, k_n).$$

This is a finite sum and thus well-defined since $\Delta_{k_i}^{d+1} G(k_1, \dots, k_n) = 0$ if $\deg_{k_i} G(k_1, \dots, k_n) = d$.

Lemma 1. *Let $p(X_1, \dots, X_n)$ be a formal power series in $(X_1, \dots, X_n)$ over $\mathbb{C}$ with non-zero constant term. Then $p(\Delta_{k_1}, \dots, \Delta_{k_n})$ is invertible as an operator over $\mathbb{C}[k_1, \dots, k_n]$, i.e. there exists a formal power series $q(X_1, \dots, X_n)$ with*

$$\begin{aligned} p(\Delta_{k_1}, \dots, \Delta_{k_n}) q(\Delta_{k_1}, \dots, \Delta_{k_n}) F(k_1, \dots, k_n) \\ = q(\Delta_{k_1}, \dots, \Delta_{k_n}) p(\Delta_{k_1}, \dots, \Delta_{k_n}) F(k_1, \dots, k_n) = F(k_1, \dots, k_n) \end{aligned}$$

for all polynomials $F(k_1, \dots, k_n)$. Moreover

$$\begin{aligned} \deg_{k_{i_1}, \dots, k_{i_m}} G(k_1, \dots, k_n) = \deg_{k_{i_1}, \dots, k_{i_m}} p(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n) = \\ \deg_{k_{i_1}, \dots, k_{i_m}} q(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n) \end{aligned}$$

for all $(i_1, i_2, \dots, i_m)$ with $1 \leq i_1 < i_2 < \dots < i_m \leq n$, where $\deg_{k_{i_1}, \dots, k_{i_m}} G(k_1, \dots, k_n)$ denotes the degree of $G(k_1, \dots, k_n)$ as a polynomial in $(k_{i_1}, \dots, k_{i_m})$.

Proof. The assertion follows from the fact that $p(X_1, \dots, X_n)$ is invertible in the (commutative) algebra of formal power series over $\mathbb{C}$ if (and only if) $p(X_1, \dots, X_n)$ has a non-zero constant term. This is because $p(X_1, \dots, X_n) q(X_1, \dots, X_n) = 1$ is

equivalent to $a_{0,\dots,0}b_{0,\dots,0} = 1$ and

$$\sum_{\substack{(i_1,\dots,i_n),(j_1,\dots,j_n) \in (\mathbb{Z}^{\geq 0})^n \\ (i_1,\dots,i_n)+(j_1,\dots,j_n)=(r_1,\dots,r_n)}} a_{i_1,\dots,i_n} b_{j_1,\dots,j_n} = 0$$

for $(r_1, \dots, r_n) \neq (0, \dots, 0)$, where

$$p(X_1, \dots, X_n) = \sum_{(j_1,\dots,j_n) \in (\mathbb{Z}^{\geq 0})^n} a_{j_1,\dots,j_n} X_1^{j_1} \cdots X_n^{j_n}.$$

and

$$q(X_1, \dots, X_n) = \sum_{(j_1,\dots,j_n) \in (\mathbb{Z}^{\geq 0})^n} b_{j_1,\dots,j_n} X_1^{j_1} \cdots X_n^{j_n}.$$

By assumption $a_{0,\dots,0} \neq 0$ and, consequently, the equations allow us to determine the coefficients $b_{r_1,\dots,r_n}$ by induction with respect to $r_1 + \dots + r_n$. The assertion about the degree follows from the fact that

$$\begin{aligned} \deg_{k_{i_1},\dots,k_{i_m}} p(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n) &\leq \deg_{k_{i_1},\dots,k_{i_m}} G(k_1, \dots, k_n) \\ &= \deg_{k_{i_1},\dots,k_{i_m}} q(\Delta_{k_1}, \dots, \Delta_{k_n}) p(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n) \\ &\leq \deg_{k_{i_1},\dots,k_{i_m}} p(\Delta_{k_1}, \dots, \Delta_{k_n}) G(k_1, \dots, k_n). \quad \square \end{aligned}$$

We define $V_{x,y} = (\text{id} + E_y \Delta_x) = E_x + \Delta_x \Delta_y$. In the following lemma we will see why this operator is of significance for the recursion underlying $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$. (The lemma is equivalent to [2, Lemma 2].) It will be used for showing that the degree of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is no greater than $n - 1$ in every k_i .

Lemma 2. *Let $a(x, y)$ be a polynomial in x and y which is of degree no greater than R in each of x and y . Moreover, assume that $(\text{id} + S_{x,y})V_{x,y}a(x, y)$ is of degree no greater than R as a polynomial in x and y , i.e. a linear combination of monomials $x^m y^n$ with $m + n \leq R$. Then*

$$\sum_{(x,y)}^{(k_1,k_2,k_3)} a(x, y) = \sum_{x=k_1}^{k_2} \sum_{y=k_2}^{k_3} a(x, y) - a(k_2, k_2) \quad (3.1)$$

is of degree no greater than $R + 2$ in k_2 . Moreover, if $(\text{id} + S_{x,y})V_{x,y}a(x, y) = 0$ then the degree of (3.1) in k_2 is no greater than $R + 1$.

Proof. First note that, by Lemma 1, $V_{x,y} + V_{y,x} = 2 \text{id} + \Delta_x + \Delta_y + 2\Delta_x \Delta_y$ is invertible. Thus

$$(\text{id} + S_{x,y}) \frac{V_{x,y}}{V_{x,y} + V_{y,x}} + \frac{V_{y,x}}{V_{x,y} + V_{y,x}} (\text{id} - S_{x,y}) = \text{id}, \quad (3.2)$$

since $V_{x,y} + V_{y,x}$ and $S_{x,y}$ commute. Moreover,

$$(\text{id} + S_{x,y}) \frac{V_{x,y}}{V_{x,y} + V_{y,x}} a(x, y) = \frac{1}{V_{x,y} + V_{y,x}} (\text{id} + S_{x,y}) V_{x,y} a(x, y).$$

By Lemma 1, the degree of this expression in x and y is equal to the degree of $(\text{id} + S_{x,y})V_{x,y}a(x, y)$ in x and y and the expression vanishes if and only if $(\text{id} + S_{x,y})V_{x,y}a(x, y)$ vanishes. Thus, by (3.2), it suffices to show that the degree of

$$\sum_{(x,y)}^{(k_1,k_2,k_3)} \frac{V_{y,x}}{V_{x,y} + V_{y,x}} (\text{id} - S_{x,y})a(x, y)$$

in k_2 is no greater than $R + 1$. Once more the degree estimation from Lemma 1 implies that this can be reduced to showing the following. If we define $b(x, y) = V_{y,x}(\text{id} - S_{x,y})\binom{x}{p}\binom{y}{q}$ then the degree of $\sum_{(x,y)}^{(k_1,k_2,k_3)} b(x, y)$ in k_2 is no greater than $\max(p, q) +$

1. In order to do so, observe that

$$\begin{aligned} b(x, y) &= V_{y,x}(\text{id} - S_{x,y})\binom{x}{p}\binom{y}{q} \\ &= \binom{x}{p}\binom{y}{q} - \binom{x}{q}\binom{y}{p} + \binom{x+1}{p}\binom{y}{q-1} - \binom{x+1}{q}\binom{y}{p-1}. \end{aligned}$$

Therefore, and by the summation formula

$$\sum_{x=a}^b \binom{x}{n} = \sum_{x=a}^b \binom{x+1}{n+1} - \binom{x}{n+1} = \binom{b+1}{n+1} - \binom{a}{n+1},$$

we have

$$\begin{aligned} \sum_{x=k_1}^{k_2} \sum_{y=k_2}^{k_3} b(x, y) - b(k_2, k_2) &= \left(\binom{k_2+1}{p+1} - \binom{k_1}{p+1} \right) \left(\binom{k_3+1}{q+1} - \binom{k_2}{q+1} \right) \\ &\quad - \left(\binom{k_2+1}{q+1} - \binom{k_1}{q+1} \right) \left(\binom{k_3+1}{p+1} - \binom{k_2}{p+1} \right) \\ &\quad + \left(\binom{k_2+2}{p+1} - \binom{k_1+1}{p+1} \right) \left(\binom{k_3+1}{q} - \binom{k_2}{q} \right) \\ &\quad - \left(\binom{k_2+2}{q+1} - \binom{k_1+1}{q+1} \right) \left(\binom{k_3+1}{p} - \binom{k_2}{p} \right) \\ &\quad - \binom{k_2+1}{p} \binom{k_2}{q-1} + \binom{k_2+1}{q} \binom{k_2}{p-1}. \end{aligned}$$

If we repeatedly apply the identity

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

to this expression, we finally see that this is a polynomial in k_2 of degree no greater than $\max(p, q) + 1$. $\square$

In order to use this lemma to compute the degree of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ in every k_i , we need to show that $(\text{id} + S_{k_i, k_{i+1}})V_{k_i, k_{i+1}}\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil}) = 0$ for all i . This

will be a consequence of the following lemma, which implies that

$$(\text{id} + S_{k_i, k_{i+1}})V_{k_i, k_{i+1}} \left(\sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} a(l_1, \dots, l_{m-1}) \right)$$

is expressible as a certain sum of

$$(\text{id} + S_{l_{i-1}, l_i})V_{l_{i-1}, l_i} a(l_1, \dots, l_{m-1})$$

and

$$(\text{id} + S_{l_i, l_{i+1}})V_{l_i, l_{i+1}} a(l_1, \dots, l_{m-1}).$$

It is yet another result, which manifests the connection of $V_{x,y}$ and the recursion. The lemma is Lemma 3 of [2] and we omit its proof here. In order to simplify the statement we use the following notation $T_{x,y} = (\text{id} + S_{x,y})V_{x,y}$.

Lemma 3. *Let $f(l_1, l_2, l_3)$ be a function on $\mathbb{Z}^3$ with values in $\mathbb{C}$ and define*

$$g(k_1, k_2, k_3, k_4) = \sum_{(l_1, l_2, l_3)}^{(k_1, k_2, k_3, k_4)} f(l_1, l_2, l_3).$$

Then

$$\begin{aligned} & T_{k_2, k_3} g(k_1, k_2, k_3, k_4) \\ &= -\frac{1}{2} \left(\sum_{l_1=k_2+1}^{k_3} \sum_{l_2=k_2+1}^{k_3} \sum_{l_3=k_2}^{k_4} T_{l_1, l_2} f(l_1, l_2, l_3) + \sum_{l_1=k_1}^{k_2+1} \sum_{l_2=k_2}^{k_3-1} \sum_{l_3=k_2}^{k_3-1} T_{l_2, l_3} f(l_1, l_2, l_3) \right) \\ &+ \frac{1}{2} \left(\sum_{l_1=k_2}^{k_3-1} \sum_{l_2=k_2}^{k_3-1} \Delta_{l_2} (\text{id} + E_{l_1}) T_{l_1, l_2} f(l_1, l_2, k_2) - \sum_{l_2=k_2}^{k_3-1} \sum_{l_3=k_2}^{k_3-1} \Delta_{l_2} (\text{id} + E_{l_3}) T_{l_2, l_3} f(k_2 + 1, l_2, l_3) \right) \\ &+ \frac{1}{2} \left(T_{l_1, l_2} f(l_1, l_2, k_2 + 1)|_{(l_1, l_2)=(k_2, k_2)} - T_{l_1, l_2} f(l_1, l_2, k_3 + 1)|_{(l_1, l_2)=(k_2, k_2)} \right. \\ &\quad \left. + T_{l_2, l_3} f(k_2, l_2, l_3)|_{(l_2, l_3)=(k_2, k_2)} - T_{l_2, l_3} f(k_3, l_2, l_3)|_{(l_2, l_3)=(k_2, k_2)} \right. \\ &\quad \left. - T_{l_1, l_2} f(l_1, l_2, k_2 + 1)|_{(l_1, l_2)=(k_2, k_3)} - T_{l_2, l_3} f(k_2, l_2, l_3)|_{(l_2, l_3)=(k_2, k_3)} \right). \end{aligned}$$

Moreover, for a function $h(l_1, l_2)$ on $\mathbb{Z}^2$,

$$T_{k_1, k_2} \sum_{(l_1, l_2)}^{(k_1, k_2, k_3)} h(l_1, l_2) = -\frac{1}{2} \sum_{l_1=k_1}^{k_2-1} \sum_{l_2=k_1}^{k_2-1} T_{l_1, l_2} h(l_1, l_2).$$

This proves the assertion preceding the lemma for $m = 3, 4$. For $m = 2$ observe that

$$\begin{aligned} (\text{id} + S_{k_1, k_2})V_{k_1, k_2} \sum_{(l_1)}^{(k_1, k_2)} a(l_1) &= (\text{id} + S_{k_1, k_2}) \left(\sum_{l_1=k_1}^{k_2} a(l_1) - a(k_1) \right) \\ &= \sum_{l_1=k_1+1}^{k_2} a(l_1) + \sum_{l_2=k_2+1}^{k_1} a(l_1) = 0. \end{aligned}$$

In order to use Lemma 3 to prove the assertion for $m \geq 5$, we need a merging rule for (2.1). Let $f(x, z)$ be a function on $\mathbb{Z}^2$. Then the operator $I_{x,z}^y$ is defined as follows.

$$I_{x,z}^y f(x, z) = f(y-1, y) + f(y, y+1) - f(y-1, y+1) = V_{x,z} f(x, z)|_{(x=y-1, z=y)}$$

Using this operator, we have

$$\sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} a(l_1, \dots, l_{m-1}) = I_{w,x}^{k_{i-1}} I_{y,z}^{k_{i+2}} \sum_{(l_1, \dots, l_{i-2})}^{(k_1, \dots, k_{i-2}, w)} \sum_{(l_{i-1}, l_i, l_{i+1})}^{(x, k_i, k_{i+1}, y)} \sum_{(l_{i+2}, \dots, l_n)}^{(z, k_{i+3}, \dots, l_{m-1})} a(l_1, \dots, l_{m-1}) \quad (3.3)$$

and this enables one to prove the assertion for $m \geq 5$. (For details see [2, Section 4].)

After noting that $(\text{id} + S_{x,y})g(x, y) = 0$ if and only if $g(x, y)$ is antisymmetric in x and y , we finally obtain the following.

Corollary 1. *Suppose $a(l_1, \dots, l_{m-1})$ is a function on $\mathbb{Z}^{m-1}$ such that $V_{l_i, l_{i+1}} a(l_1, \dots, l_{m-1})$ is antisymmetric in l_i and l_{i+1} for all i . Then*

$$V_{k_i, k_{i+1}} \sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)} a(l_1, \dots, l_{m-1})$$

is antisymmetric in k_i and k_{i+1} for all i .

4. CHARACTERIZING PROPERTIES OF $\gamma(n; k_1, \dots, k_{\lceil n/2 \rceil})$

We apply the results from the previous section to $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$: Corollary 1 implies by induction with respect to n that $V_{k_i, k_{i+1}} \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is antisymmetric in k_i and k_{i+1} for all i . Lemma 2 and the merging rule (3.3) then implies by induction with respect to n that the degree of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ in k_i is no greater than $n-1$. We summarize this in the following lemma.

Lemma 4. *For $n \geq 1$, $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is a polynomial of degree no greater than $n-1$ in every k_i . Furthermore, $V_{k_i, k_{i+1}} \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is antisymmetric in k_i and k_{i+1} for all i .*

It will be shown that the properties from the previous lemma together with the property in the following lemma characterize γ up to a multiplicative rational constant.

Lemma 5. *If n is even then*

$$\gamma(n, x; k_1, \dots, k_{n/2-1}, k_{n/2}) = -\gamma(n, x; k_1, \dots, k_{n/2-1}, 2x+2-k_{n/2})$$

and if n is odd then

$$\gamma(n, x; k_1, \dots, k_{(n-1)/2}, k_{(n+1)/2}) = \gamma(n, x; k_1, \dots, k_{(n-1)/2}, 2x+1-k_{(n+1)/2}).$$

In order to prove this lemma, we need another lemma.

Lemma 6. (1) *Let $f(l_1)$ be such that $f(l_1) = -f(2x+2-l_1)$ for all l_1 and define*

$$g(k_1, k_2) = \sum_{l_1=k_1}^{k_2} f(l_1). \text{ Then } g(k_1, k_2) = g(k_1, 2x+1-k_2) \text{ for all } k_1, k_2.$$

(2) *Let $f(l_1, l_2)$ be such that $f(l_1, l_2) = f(l_1, 2x+1-l_2)$ for all l_1, l_2 and*

$$(\text{id} + S_{l_1, l_2}) V_{l_1, l_2} f(l_1, l_2) = 0.$$

Define

$$g(k_1, k_2) = \sum_{(l_1, l_2)}^{(k_1, k_2, x)} f(l_1, l_2) = \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) - f(k_2, k_2).$$

Then $g(k_1, k_2) = -g(k_1, 2x+2-k_2)$ for all k_1, k_2 .

Proof of Lemma 6. (1) By definition,

$$g(k_1, 2x+1-k_2) = \sum_{l_1=k_1}^{2x+1-k_2} f(l_1) = \sum_{l_1=k_1}^{k_2} f(l_1) + \sum_{l_1=k_2+1}^{2x+1-k_2} f(l_1).$$

The assertion follows since $\sum_{l_1=k_2+1}^{2x+1-k_2} f(l_1) = 0$. This is because

$$\sum_{l_1=k_2+1}^{2x+1-k_2} f(l_1) = \sum_{l_1=k_2+1}^{2x+1-k_2} -f(2x+2-l_1) = - \sum_{l_1=k_2+1}^{2x+1-k_2} f(l_1).$$

(2) Observe that

$$\begin{aligned} g(k_1, 2x+2-k_2) &= \sum_{l_1=k_1}^{2x+2-k_2} \sum_{l_2=2x+2-k_2}^x f(l_1, l_2) - f(2x+2-k_2, 2x+2-k_2) \\ &= \sum_{l_1=k_1}^{k_2} \sum_{l_2=2x+2-k_2}^x f(l_1, l_2) + \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=2x+2-k_2}^x f(l_1, l_2) - f(2x+2-k_2, 2x+2-k_2) \\ &= \sum_{l_1=k_1}^{k_2} \sum_{l_2=2x+2-k_2}^x f(l_1, 2x+1-l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=x+1}^{2x+1-k_2} f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=x+1}^{2x+1-k_2} f(l_1, l_2) \\ &\quad - f(2x+2-k_2, 2x+2-k_2) \\ &= \sum_{l_1=k_1}^{k_2} \sum_{l_2=x+1}^{k_2-1} f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=x+1}^{2x+1-k_2} f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=x+1}^{2x+1-k_2} f(l_1, 2x+1-l_2) \\ &\quad - f(2x+2-k_2, 2x+2-k_2) \\ &= - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=x+1}^{2x+1-k_2} f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^x f(l_1, l_2) \\ &\quad - f(2x+2-k_2, 2x+2-k_2) \\ &= - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) - \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^{2x+1-k_2} f(l_1, l_2) - f(2x+2-k_2, 2x+2-k_2). \quad (4.1) \end{aligned}$$

Moreover, we have $E_x^{-1}(\text{id} + S_{x,y})V_{x,y} = (\text{id} + E_y E_x^{-1} S_{x,y})(\text{id} + E_x^{-1} \Delta_x \Delta_y)$, and, therefore, $(\text{id} + E_{l_2} E_{l_1}^{-1} S_{l_1, l_2})(\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2})f(l_1, l_2) = 0$. This implies that

$$\sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^{2x+1-k_2} (\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2})f(l_1, l_2) = 0,$$

since

$$\begin{aligned}
& \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^{2x+1-k_2} (\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2}) f(l_1, l_2) \\
&= \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^{2x+1-k_2} -E_{l_2} E_{l_1}^{-1} S_{l_1, l_2} (\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2}) f(l_1, l_2) \\
&= - \sum_{l_1=k_2}^{2x+1-k_2} \sum_{l_2=k_2+1}^{2x+2-k_2} S_{l_1, l_2} (\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2}) f(l_1, l_2) \\
&= - \sum_{l_2=k_2}^{2x+1-k_2} \sum_{l_1=k_2+1}^{2x+2-k_2} (\text{id} + E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2}) f(l_1, l_2).
\end{aligned}$$

Thus, (4.1) is equal to

$$\begin{aligned}
& - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) + \frac{1}{2} \sum_{l_1=k_2+1}^{2x+2-k_2} \sum_{l_2=k_2}^{2x+1-k_2} E_{l_1}^{-1} \Delta_{l_1} \Delta_{l_2} f(l_1, l_2) - f(2x+2-k_2, 2x+2-k_2) \\
&= - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) - \frac{1}{2} f(2x+2-k_2, 2x+2-k_2) + \frac{1}{2} f(k_2, k_2) - \frac{1}{2} f(2x+2-k_2, k_2) \\
&\quad - \frac{1}{2} f(k_2, 2x+2-k_2). \quad (4.2)
\end{aligned}$$

Next observe that

$$\begin{aligned}
0 &= ((\text{id} + S_{l_1, l_2}) V_{l_1, l_2} f(l_1, l_2))|_{(l_1, l_2) = (k_2-1, 2x+1-k_2)} \\
&= f(k_2-1, 2x+1-k_2) + f(k_2, 2x+2-k_2) - f(k_2-1, 2x+2-k_2) \\
&\quad + f(2x+1-k_2, k_2-1) + f(2x+2-k_2, k_2) - f(2x+1-k_2, k_2).
\end{aligned}$$

We replace $-\frac{1}{2} f(2x+2-k_2, k_2) - \frac{1}{2} f(k_2, 2x+2-k_2)$ in (4.2) by

$$\frac{1}{2} f(k_2-1, 2x+1-k_2) + \frac{1}{2} f(2x+1-k_2, k_2-1) - \frac{1}{2} f(k_2-1, 2x+2-k_2) - \frac{1}{2} f(2x+1-k_2, k_2)$$

and, consequently, (4.2) is equal to

$$\begin{aligned}
& - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) + f(k_2, k_2) - \frac{1}{2} (f(2x+2-k_2, 2x+2-k_2) + f(k_2, k_2) \\
& - f(k_2-1, 2x+1-k_2) - f(2x+1-k_2, k_2-1) + f(k_2-1, 2x+2-k_2) + f(2x+1-k_2, k_2)) \\
&= - \sum_{l_1=k_1}^{k_2} \sum_{l_2=k_2}^x f(l_1, l_2) + f(k_2, k_2) - \frac{1}{2} (f(2x+2-k_2, 2x+2-k_2) + f(k_2, k_2) \\
& - f(k_2-1, k_2) - f(2x+1-k_2, 2x+2-k_2) + f(k_2-1, k_2-1) + f(2x+1-k_2, 2x+1-k_2)). \quad (4.3)
\end{aligned}$$

Finally,

$$0 = ((\text{id} + S_{l_1, l_2})V_{l_1, l_2}f(l_1, l_2))|_{(l_1, l_2)=(l, l)} = 2(f(l, l) + f(l+1, l+1) - f(l, l+1))$$

implies that (4.3) is equal to $-g(k_1, k_2)$. $\square$

Proof of Lemma 5. We use induction with respect to n . For $n = 2$ the assertion is easy to check. We assume that $n \geq 3$ and first consider the case that n is odd. By the recursion, it suffices to show that

$$\begin{aligned} & \sum_{(l_1, \dots, l_{(n-1)/2})}^{(k_1, \dots, k_{(n-1)/2}, k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}) \\ &= \sum_{(l_1, \dots, l_{(n-1)/2})}^{(k_1, \dots, k_{(n-1)/2}, 2x+1-k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}) \quad (4.4) \end{aligned}$$

By the induction hypothesis and by Lemma 6 (1) we know that

$$\sum_{(l_{(n-1)/2})}^{(k''_{(n-1)/2}, k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}) = \sum_{(l_{(n-1)/2})}^{(k''_{(n-1)/2}, 2x+1-k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}).$$

The assertion follows, since the left hand side of (4.4) is equal to

$$I_{k_{(n-1)/2}}^{k'_{(n-1)/2}, k''_{(n-1)/2}} \sum_{(l_1, \dots, l_{(n-3)/2})}^{(k_1, \dots, k'_{(n-1)/2})} \sum_{(l_{(n-1)/2})}^{(k''_{(n-1)/2}, k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2})$$

and the right hand side of (4.4) is equal to

$$I_{k_{(n-1)/2}}^{k'_{(n-1)/2}, k''_{(n-1)/2}} \sum_{(l_1, \dots, l_{(n-3)/2})}^{(k_1, \dots, k'_{(n-1)/2})} \sum_{(l_{(n-1)/2})}^{(k''_{(n-1)/2}, 2x+1-k_{(n+1)/2})} \gamma(n-1, x; l_1, \dots, l_{(n-1)/2}).$$

In the case that n is even, Lemma 6 (2) is used in a similar way. $\square$

5. DERIVATION OF THE OPERATOR FORMULA

By Lemma 4, we know that $V_{k_i, k_{i+1}}\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is antisymmetric in k_i and k_{i+1} for all i . Although the operators $V_{k_i, k_{i+1}}$ commute, this clearly does not imply that

$$V_{k_1, k_2} V_{k_2, k_3} \cdots V_{k_{\lceil n/2 \rceil - 1}, k_{\lceil n/2 \rceil}} \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

is antisymmetric in $(k_1, k_2, \dots, k_{\lceil n/2 \rceil})$. However, it is not hard to see that

$$\left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} V_{k_p, k_q} \right) \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil}) \quad (5.1)$$

is antisymmetric in $(k_1, k_2, \dots, k_{\lceil n/2 \rceil})$. This is a consequence of the following lemma, which generalizes [2, Lemma 4]. The proof is analogous to the proof of [2, Lemma 4] and thus we omit it here.

Lemma 7. *Let $W_{x,y}$ be an operator in x and y , which is invertible as operator over $\mathbb{C}[x, y]$, and $W_{x_1, y_1} W_{x_2, y_2} = W_{x_2, y_2} W_{x_1, y_1}$ for all x_1, x_2, y_1, y_2 . Moreover, let $a(k_1, \dots, k_m)$ be a polynomial in $(k_1, \dots, k_m)$. Then $W_{k_i, k_{i+1}} a(k_1, \dots, k_m)$ is antisymmetric in k_i and k_{i+1} for all i if and only if*

$$\left(\prod_{1 \leq p < q \leq m} W_{k_p, k_q} \right) a(k_1, \dots, k_m)$$

is antisymmetric in $(k_1, \dots, k_m)$.

We denote the polynomials in (5.1) by $\gamma^*(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ and list them for $n = 1, 2, 3, 4, 5$.

$$\begin{aligned} & 1, x - k_1 + 1, \frac{1}{2}(2x + 1 - k_1 - k_2)(k_2 - k_1), \frac{1}{6}(k_2 - k_1)(2x^3 - 3k_1x^2 - 3k_2x^2 + 6x^2 + k_1^2x + k_2^2x \\ & - 6k_1x + 4k_1k_2x - 6k_2x + 12x + k_1^2 - k_1k_2^2 + k_2^2 - 6k_1 - k_1^2k_2 + 4k_1k_2 - 6k_2 + 11), \\ & \frac{1}{48}(k_2 - k_1)(k_3 - k_1)(k_3 - k_2)(8x^3 - 8k_1x^2 - 8k_2x^2 - 8k_3x^2 + 12x^2 + 2k_1^2x + 2k_2^2x + 2k_3^2x - 8k_1x \\ & + 6k_1k_2x - 8k_2x + 6k_1k_3x + 6k_2k_3x - 8k_3x + 30x + k_1^2 - k_1k_2^2 + k_2^2 - k_1k_3^2 - k_2k_3^2 + k_3^2 - 10k_1 \\ & - k_1^2k_2 + 3k_1k_2 - 10k_2 - k_1^2k_3 - k_2^2k_3 + 3k_1k_3 - 2k_1k_2k_3 + 3k_2k_3 - 10k_3 + 25) \end{aligned}$$

Although this list is shorter than the analog list for $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ (this is due to the factor $\prod_{1 \leq i < j \leq \lceil n/2 \rceil} (k_j - k_i)$, which is a consequence of the antisymmetry of the

polynomial), it is still hard to guess the general pattern of γ^* . Thus we will apply a further operator to γ^* , in order to obtain a polynomial which factorizes into linear factors over $\mathbb{Q}$ and for which it is easy to recognize a pattern. This operator will have the property that it does not destroy the antisymmetry of the polynomial but restores the symmetry property of γ given in Lemma 5. In the end, the fact that our operators are invertible will allow us to “divide” and give a formula for γ itself.

The next lemma shows that the application of an operator, which is a symmetric polynomial in the shift operators, to an antisymmetric polynomial retains the antisymmetry.

Lemma 8. *Let $a(k_1, \dots, k_m)$ be a polynomial that is antisymmetric in $(k_1, \dots, k_m)$ and $p(X_1, \dots, X_m)$ be a polynomial in $X_1, X_1^{-1}, X_2, X_2^{-1}, \dots, X_m, X_m^{-1}$, which is symmetric in $(X_1, \dots, X_m)$. Then $p(E_{k_1}, \dots, E_{k_m})a(k_1, \dots, k_m)$ is antisymmetric in $(k_1, \dots, k_m)$.*

Proof. Let $\sigma \in \mathcal{S}_m$ be a permutation and $p(X_1, \dots, X_m) = \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} X_1^{i_1} \cdots X_m^{i_m}$. The symmetry of $p(X_1, \dots, X_m)$ implies that $c_{i_1, \dots, i_m} = c_{i_{\sigma(1)}, \dots, i_{\sigma(m)}}$. Thus

$$\begin{aligned}
p(E_{k_1}, \dots, E_{k_m})a(k_1, \dots, k_m) &= \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} E_{k_1}^{i_1} \cdots E_{k_m}^{i_m} a(k_1, \dots, k_m) \\
&= \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_1 + i_1, \dots, k_m + i_m) \\
&= \text{sgn } \sigma \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_{\sigma(1)} + i_{\sigma(1)}, \dots, k_{\sigma(m)} + i_{\sigma(m)}) \\
&= \text{sgn } \sigma \sum_{(i_1, \dots, i_m)} c_{i_{\sigma(1)}, \dots, i_{\sigma(m)}} a(k_{\sigma(1)} + i_{\sigma(1)}, \dots, k_{\sigma(m)} + i_{\sigma(m)}) \\
&= \text{sgn } \sigma \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_{\sigma(1)} + i_1, \dots, k_{\sigma(m)} + i_m) \\
&= \text{sgn } \sigma (p(E_{l_1}, \dots, E_{l_m})a(l_1, \dots, l_m))|_{(l_1, \dots, l_m) = (k_{\sigma(1)}, \dots, k_{\sigma(m)})}. \quad \square
\end{aligned}$$

In the following lemma we identify operators whose application do not destroy symmetry properties of the type given in Lemma 5.

Lemma 9. *Let $a(k_1, \dots, k_m)$ be a polynomial such that*

$$a(k_1, \dots, k_m) = \sigma \cdot a(k_1, \dots, k_{m-1}, d - k_m)$$

for $\sigma, d \in \mathbb{R}$ and $p(X_1, \dots, X_m)$ be a polynomial in $X_1, X_1^{-1}, X_2, X_2^{-1}, \dots, X_m, X_m^{-1}$ such that

$$p(X_1, \dots, X_m) = p(X_1, \dots, X_{m-1}, X_m^{-1}).$$

Set $b(k_1, \dots, k_m) = p(E_{k_1}, \dots, E_{k_m})a(k_1, \dots, k_m)$. Then we have

$$b(k_1, \dots, k_m) = \sigma \cdot b(k_1, \dots, k_{m-1}, d - k_m)$$

as well.

Proof. Let $p(X_1, \dots, X_m) = \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} X_1^{i_1} \cdots X_m^{i_m}$. By assumption $c_{i_1, \dots, i_{m-1}, i_m} = c_{i_1, \dots, i_{m-1}, -i_m}$. Therefore,

$$\begin{aligned}
p(E_{k_1}, \dots, E_{k_m})a(k_1, \dots, k_m) &= \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_1 + i_1, \dots, k_m + i_m) \\
&= \sigma \cdot \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_1 + i_1, \dots, k_{m-1} + i_{m-1}, d - k_m - i_m) \\
&= \sigma \cdot \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_{m-1}, -i_m} a(k_1 + i_1, \dots, k_{m-1} + i_{m-1}, d - k_m - i_m) \\
&= \sigma \cdot \sum_{(i_1, \dots, i_m)} c_{i_1, \dots, i_m} a(k_1 + i_1, \dots, k_{m-1} + i_{m-1}, d - k_m + i_m) \\
&= \sigma \cdot (p(E_{l_1}, \dots, E_{l_m})a(l_1, \dots, l_m))|_{(l_1, \dots, l_m) = (k_1, \dots, k_{m-1}, d - k_m)}. \quad \square
\end{aligned}$$

The previous two lemmas suggest to look for an operator $p(E_{k_1}, \dots, E_{k_{\lceil n/2 \rceil}})$, which is, on the one hand, symmetric in $(k_1, \dots, k_{\lceil n/2 \rceil})$ and, on the other hand, has the property that the composition

$$p(E_{k_1}, \dots, E_{k_{\lceil n/2 \rceil}}) \left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (\text{id} + E_{k_q} \Delta_{k_p}) \right)$$

is invariant under the replacement of $E_{k_{\lceil n/2 \rceil}}$ by $E_{k_{\lceil n/2 \rceil}}^{-1}$. This is accomplished in the following lemma.

Lemma 10. *The polynomial*

$$\left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (\text{id} + E_{k_q} \Delta_{k_p}) E_{k_p}^{-1} (\text{id} + E_{k_q}^{-1} \Delta_{k_p}) \right) \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

is antisymmetric in $(k_1, \dots, k_{\lceil n/2 \rceil})$. Moreover, if n is odd then the polynomial is invariant under the replacement of k_i by $2x + 1 - k_i$ and if n is even then the replacement of k_i by $2x + 2 - k_i$ only changes the sign of the polynomial.

Proof. By Lemma 7,

$$\left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (\text{id} + E_{k_q} \Delta_{k_p}) \right) \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

is antisymmetric in $(k_1, \dots, k_{\lceil n/2 \rceil})$. Lemma 8 and the fact that

$$\prod_{1 \leq p < q \leq \lceil n/2 \rceil} X_p^{-1} (1 + X_q^{-1} (X_p - 1)) = \prod_{1 \leq p < q \leq \lceil n/2 \rceil} (X_p^{-1} + X_q^{-1} - X_p^{-1} X_q^{-1})$$

is symmetric in $(X_1, \dots, X_{\lceil n/2 \rceil})$ imply that the expression in the statement of the lemma is still antisymmetric in $(k_1, \dots, k_{\lceil n/2 \rceil})$.

Next observe that the operator in the statement of the lemma is a polynomial in the shift operators $E_{k_i}^{\pm 1}$, which is invariant under the replacement of $E_{k_{\lceil n/2 \rceil}}$ by $E_{k_{\lceil n/2 \rceil}}^{-1}$. Therefore, by Lemma 5 and by Lemma 9, the second assertion in lemma follows for $i = \lceil n/2 \rceil$. The assertion for general i follows from the antisymmetry of the polynomial. $\square$

The next lemma shows that the previous lemma together with the degree estimation (Lemma 4) determines $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ up to multiplicative constant, which only depends on n .

Lemma 11. *Let $p(k_1, \dots, k_{\lceil n/2 \rceil})$ be an antisymmetric polynomial in $(k_1, \dots, k_{\lceil n/2 \rceil})$ over $\mathbb{C}$ of degree no greater than $n - 1$ in every k_i which is, in the case that n is odd, invariant under the replacement of k_i by $2x + 1 - k_i$ for every i and, in the case that n is even, has the property that the replacement of k_i by $2x + 2 - k_i$ only changes the sign of the polynomial. Then $p(k_1, \dots, k_{\lceil n/2 \rceil})$ equals*

$$C \cdot \prod_{1 \leq i < j \leq (n+1)/2} (k_j - k_i)(2x + 1 - k_i - k_j)$$

if n is odd and

$$C \cdot \left(\prod_{1 \leq i < j \leq n/2} (k_j - k_i)(2x + 2 - k_i - k_j) \right) \prod_{i=1}^{n/2} (x + 1 - k_i)$$

if n is even, where C is a constant in $\mathbb{C}$.

Proof. We only consider the case that n is even for the other case is analogous. A polynomial $p(k_1, \dots, k_{n/2})$ that is antisymmetric in $(k_1, \dots, k_{n/2})$ must have $k_j - k_i$ as a factor since

$$p(k_1, \dots, k_{i-1}, k_i, k_{i+1}, \dots, k_{j-1}, k_i, k_{j+1}, \dots, k_{n/2}) = -p(k_1, \dots, k_{i-1}, k_i, k_{i+1}, \dots, k_{j-1}, k_i, k_{j+1}, \dots, k_{n/2}).$$

This is because the polynomial changes the sign if we exchange the element in the i -th position with the element in the j -th position. If it furthermore has the property that it will change the sign if k_j is replaced by $2x + 2 - k_j$ then the polynomial has a zero at $k_j = 2x + 2 - k_i$ which explains the factor $2x + 2 - k_i - k_j$. Moreover it has a zero at $k_i = x + 1$ for every i , since

$$p(k_1, \dots, k_{i-1}, x + 1, k_{i+1}, \dots, k_{n/2}) = -p(k_1, \dots, k_{i-1}, x + 1, k_{i+1}, \dots, k_{n/2}),$$

which follows from $2x + 2 - (x + 1) = x + 1$. $\square$

Consequently, by Lemma 4, Lemma 10 and Lemma 11,

$$\left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (\text{id} + E_{k_q} \Delta_{k_p}) E_{k_p}^{-1} (\text{id} + E_{k_q}^{-1} \Delta_{k_p}) \right) \gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

is equal to the polynomials given in Lemma 11. This determines $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ up to a multiplicative complex constant C_n . This is because the operators $(\text{id} + E_{k_q} \Delta_{k_p})$ and $(\text{id} + E_{k_q}^{-1} \Delta_{k_p})$ are invertible by Lemma 1. In the following lemma we compute C_n .

Lemma 12. *If n is odd then*

$$C_n = \prod_{1 \leq i < j \leq (n+1)/2} \frac{1}{(j-i)(j+i-1)} = \frac{1}{(n-1)!(n-3)! \dots 2!}$$

and if n is even then

$$C_n = \prod_{1 \leq i < j \leq n/2} \frac{1}{(j-i)(j+i)} \prod_{i=1}^{n/2} \frac{1}{i} = \frac{1}{(n-1)!(n-3)! \dots 1!}.$$

Proof. We expand the polynomial $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ with respect to the basis $\prod_{i=1}^{\lceil n/2 \rceil} (k_i)_{m_i}$ and consider the coefficient of the basis element appearing in this expansion with maximal degree sequence $(m_1, m_2, \dots, m_{\lceil n/2 \rceil})$ in lexicographic order. This coefficient is equal to C_n . We show by induction with respect to n that this maximal degree sequence is $(n-1, n-3, \dots, 0)$ if n is odd and $(n-1, n-3, \dots, 1)$ if n is even.

An analysis of the definition of $\sum_{(l_1, \dots, l_{m-1})}^{(k_1, \dots, k_m)}$ and the induction hypothesis imply that this maximal basis element of $\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ is the maximal element of

$$\sum_{l_1=k_1}^{k_2-1} \sum_{l_2=k_2}^{k_3-1} \cdots \sum_{l_{(n-1)/2}=k_{(n-1)/2}}^{k_{(n+1)/2}} \prod_{i=1}^{(n-1)/2} \frac{(l_i)_{n-2i}}{(n-2i)!}$$

if n is odd and the maximal element of

$$\sum_{l_1=k_1}^{k_2-1} \sum_{l_2=k_2}^{k_3-1} \cdots \sum_{l_{n/2}=k_{n/2}}^x \prod_{i=1}^{n/2} \frac{(l_i)_{n-2i}}{(n-2i)!}$$

if n is even. □

This immediately implies the following theorem.

Theorem 2. *If n is odd then*

$$\begin{aligned} \gamma(n, x; k_1, \dots, k_{(n+1)/2}) &= \left(\prod_{1 \leq p < q \leq (n+1)/2} (\text{id} + E_{k_q} \Delta_{k_p})^{-1} E_{k_p} (\text{id} + E_{k_q}^{-1} \Delta_{k_p})^{-1} \right) \\ &\quad \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 1 - k_i - k_j)}{(j - i)(j + i - 1)} \end{aligned}$$

and if n is even then

$$\begin{aligned} \gamma(n, x; k_1, \dots, k_{n/2}) &= \left(\prod_{1 \leq p < q \leq n/2} (\text{id} + E_{k_q} \Delta_{k_p})^{-1} E_{k_p} (\text{id} + E_{k_q}^{-1} \Delta_{k_p})^{-1} \right) \\ &\quad \prod_{1 \leq i < j \leq n/2} \frac{(k_j - k_i)(2x + 2 - k_i - k_j)}{(j - i)(j + i)} \prod_{i=1}^{n/2} \frac{x + 1 - k_i}{i}. \end{aligned}$$

Proof. The assertion follows from Lemma 1, Lemma 4, Lemma 10, Lemma 11 and Lemma 12. □

We are finally able to prove the main theorem.

Proof of Theorem 1. Observe that

$$\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (1 + X_q(X_p - 1))(1 + X_q^{-1}(X_p - 1))(1 + X_q(X_p^{-1} - 1))(1 + X_q^{-1}(X_p^{-1} - 1))$$

is invariant under the replacement of X_i by X_i^{-1} . Moreover, it is symmetric in $(X_1, \dots, X_{\lceil n/2 \rceil})$, since the factor associated to the pair (p, q) is equal to

$$\begin{aligned} & (1 + X_p X_q - X_q)(1 + X_p^{-1} X_q^{-1} - X_q^{-1}) X_p X_q \\ & \quad \times (1 + X_p X_q^{-1} - X_q^{-1}) X_q \\ & \quad \times (1 + X_p^{-1} X_q - X_q) X_p \\ & \hspace{15em} \times X_p^{-2} X_q^{-2} \end{aligned}$$

and this is symmetric in X_p and X_q as every line is. Thus, by Lemma 8,

$$\begin{aligned} & \left(\prod_{1 \leq p < q \leq (n+1)/2} (\text{id} + E_{k_q} \Delta_{k_p})(\text{id} + E_{k_q}^{-1} \Delta_{k_p})(\text{id} - E_{k_q} E_{k_p}^{-1} \Delta_{k_p})(\text{id} - E_{k_q}^{-1} E_{k_p}^{-1} \Delta_{k_p}) \right) \\ & \quad \prod_{1 \leq i < j \leq (n+1)/2} (k_j - k_i)(2x + 1 - k_i - k_j) \quad (5.2) \end{aligned}$$

is antisymmetric in $(k_1, \dots, k_{(n+1)/2})$ if n is odd and

$$\begin{aligned} & \left(\prod_{1 \leq p < q \leq n/2} (\text{id} + E_{k_q} \Delta_{k_p})(\text{id} + E_{k_q}^{-1} \Delta_{k_p})(\text{id} - E_{k_q} E_{k_p}^{-1} \Delta_{k_p})(\text{id} - E_{k_q}^{-1} E_{k_p}^{-1} \Delta_{k_p}) \right) \\ & \quad \prod_{1 \leq i < j \leq n/2} (k_j - k_i)(2x + 2 - k_i - k_j) \prod_{i=1}^{n/2} (x + 1 - k_i) \quad (5.3) \end{aligned}$$

is antisymmetric in $(k_1, \dots, k_{n/2})$ if n is even. Moreover, by Lemma 9, (5.2) is invariant under the replacement of k_i by $2x + 1 - k_i$, whereas (5.3) changes the sign if k_i is replaced by $2x + 2 - k_i$. Consequently, by Lemma 11, (5.2) is equal to

$$D_n \cdot \prod_{1 \leq i < j \leq (n+1)/2} (k_j - k_i)(2x + 1 - k_i - k_j)$$

if n is odd and (5.3) is equal to

$$D_n \cdot \prod_{1 \leq i < j \leq n/2} (k_j - k_i)(2x + 2 - k_i - k_j) \prod_{i=1}^{n/2} (x + 1 - k_i)$$

if n is even. If we compare the coefficient of a monomial of maximal degree we see that $D_n = 1$. Now, if n is odd then, by Theorem 2,

$$\begin{aligned}
& \gamma(n, x; k_1, \dots, k_{(n+1)/2}) \\
&= \left(\prod_{1 \leq p < q \leq (n+1)/2} (\text{id} + E_{k_q} \Delta_{k_p})^{-1} E_{k_p} (\text{id} + E_{k_q}^{-1} \Delta_{k_p})^{-1} \right) \\
&\quad \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 1 - k_i - k_j)}{(j - i)(j + i - 1)} \\
&= \left(\prod_{1 \leq p < q \leq (n+1)/2} (\text{id} + E_{k_q} \Delta_{k_p})^{-1} E_{k_p} (\text{id} + E_{k_q}^{-1} \Delta_{k_p})^{-1} \right) \\
&\quad \left(\prod_{1 \leq p < q \leq (n+1)/2} (\text{id} + E_{k_q} \Delta_{k_p})(\text{id} + E_{k_q}^{-1} \Delta_{k_p})(\text{id} - E_{k_q} E_{k_p}^{-1} \Delta_{k_p})(\text{id} - E_{k_q}^{-1} E_{k_p}^{-1} \Delta_{k_p}) \right) \\
&\quad \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 1 - k_i - k_j)}{(j - i)(j + i - 1)} \\
&= \left(\prod_{1 \leq p < q \leq (n+1)/2} E_{k_p} (\text{id} - E_{k_q} E_{k_p}^{-1} \Delta_{k_p})(\text{id} - E_{k_q}^{-1} E_{k_p}^{-1} \Delta_{k_p}) \right) \\
&\quad \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 1 - k_i - k_j)}{(j - i)(j + i - 1)}
\end{aligned}$$

Similarly, if n is even then

$$\begin{aligned}
& \gamma(n, x; k_1, \dots, k_{n/2}) \\
&= \left(\prod_{1 \leq p < q \leq n/2} E_{k_p} (\text{id} - E_{k_q} E_{k_p}^{-1} \Delta_{k_p})(\text{id} - E_{k_q}^{-1} E_{k_p}^{-1} \Delta_{k_p}) \right) \\
&\quad \prod_{1 \leq i < j \leq n/2} \frac{(k_j - k_i)(2x + 2 - k_i - k_j)}{(j - i)(j + i)} \prod_{i=1}^{n/2} \frac{x + 1 - k_i}{i}. \quad \square
\end{aligned}$$

6. FROM OPERATOR FORMULAS TO GENERATING FUNCTIONS

In this section we follow a hint of Doron Zeilberger and translate the operator formulas into generating function results. We start out with ordinary monotone triangles. In [2] we have shown that the number of monotone triangles with bottom row $(k_1, \dots, k_n)$ is given by

$$\alpha(n; k_1, \dots, k_n) = \left(\prod_{1 \leq p < q \leq n} (\text{id} + E_{k_p} E_{k_q} - E_{k_p}) \right) \prod_{1 \leq i < j \leq n} \frac{k_j - k_i}{j - i}$$

if $k_1 < k_2 < \dots < k_n$ and $k_i \in \mathbb{Z}$. We define $\bar{\alpha}_c(n; k_1, \dots, k_n) = \prod_{1 \leq i < j \leq n} \frac{k_j - k_i}{j - i}$ if $k_l \geq c$ for all l and zero elsewhere. Then

$$\alpha(n; k_1, \dots, k_n) = \left(\prod_{1 \leq p < q \leq n} (\text{id} + E_{k_p} E_{k_q} - E_{k_p}) \right) \bar{\alpha}_c(n; k_1, \dots, k_n)$$

for all $(k_1, \dots, k_n)$ with $k_l \geq c$. We compute the generating function

$$\sum_{(k_1, \dots, k_n) \geq (-n+1, \dots, -n+1)} X_1^{k_1} \dots X_n^{k_n} \left(\prod_{1 \leq p < q \leq n} (\text{id} + E_{k_p} E_{k_q} - E_{k_p}) \right) \bar{\alpha}_0(n; k_1, \dots, k_n). \quad (6.1)$$

Thus, the coefficient of $X_1^{k_1} X_2^{k_2} \dots X_n^{k_n}$ gives the number of monotone triangles with bottom row $(k_1, \dots, k_n)$ if $0 \leq k_1 < k_2 < \dots < k_n$ and $k_i \in \mathbb{Z}$. Let

$$\prod_{1 \leq p < q \leq n} (1 + Y_p Y_q - Y_p) = \sum_{\substack{(j_1, \dots, j_n) \\ 0 \leq j_i \leq n-1}} a(j_1, \dots, j_n) Y_1^{j_1} \dots Y_n^{j_n}.$$

Using this notation, (6.1) is equal to

$$\sum_{(k_1, \dots, k_n) \geq (-n+1, \dots, -n+1)} \sum_{\substack{(j_1, \dots, j_n) \\ 0 \leq j_i \leq n-1}} a(j_1, \dots, j_n) X_1^{k_1} \dots X_n^{k_n} \bar{\alpha}_0(n; k_1 + j_1, \dots, k_n + j_n) \quad (6.2)$$

We set $(l_1, \dots, l_n) = (k_1 + j_1, \dots, k_n + j_n)$. Consequently, (6.2) is equal to

$$\sum_{\substack{(j_1, \dots, j_n) \\ 0 \leq j_i \leq n-1}} a(j_1, \dots, j_n) X_1^{-j_1} \dots X_n^{-j_n} \sum_{(l_1, \dots, l_n) \geq (-n+1+j_1, \dots, -n+1+j_n)} \bar{\alpha}_0(n; l_1, \dots, l_n) X_1^{l_1} \dots X_n^{l_n}. \quad (6.3)$$

Since $\bar{\alpha}_0(l_1, \dots, l_n) = 0$ if $l_i < 0$ for an i and $j_l \leq n-1$ for all l , (6.3) is equal to

$$\begin{aligned} & \sum_{\substack{(j_1, \dots, j_n) \\ 0 \leq j_i \leq n-1}} a(j_1, \dots, j_n) X_1^{-j_1} \dots X_n^{-j_n} \sum_{(l_1, \dots, l_n) \geq (0, \dots, 0)} \bar{\alpha}_0(n; l_1, \dots, l_n) X_1^{l_1} \dots X_n^{l_n} \\ &= \prod_{1 \leq p < q \leq n} (1 + X_p^{-1} X_q^{-1} - X_p^{-1}) \sum_{(l_1, \dots, l_n) \geq (0, \dots, 0)} X_1^{l_1} \dots X_n^{l_n} \prod_{1 \leq i < j \leq n} \frac{l_j - l_i}{j - i}. \end{aligned}$$

The Vandermonde determinant evaluation implies that

$$\prod_{1 \leq i < j \leq n} \frac{l_j - l_i}{j - i} = \det_{1 \leq i, j \leq n} \begin{pmatrix} l_i \\ j - 1 \end{pmatrix}$$

and, consequently, the generating function is equal to

$$\prod_{1 \leq p < q \leq n} (1 + X_p^{-1} X_q^{-1} - X_p^{-1}) \sum_{(l_1, \dots, l_n) \geq (0, \dots, 0)} \det_{1 \leq i, j \leq n} X_i^{l_i} \begin{pmatrix} l_i \\ j - 1 \end{pmatrix}.$$

Observe that

$$\begin{aligned} \sum_{l=0}^{\infty} X^l \binom{l}{j-1} &= \frac{X^{j-1}}{(j-1)!} \sum_{l=0}^{\infty} l(l-1) \dots (l-j+2) X^{l-j+1} = \frac{X^{j-1}}{(j-1)!} \frac{d}{dX^{j-1}} \left(\sum_{l=0}^{\infty} X^l \right) \\ &= \frac{X^{j-1}}{(j-1)!} \frac{d}{dX^{j-1}} (1-X)^{-1} = \frac{X^{j-1}}{(1-X)^j}. \end{aligned}$$

Therefore, the generating function is equal to

$$\begin{aligned} &\prod_{1 \leq p < q \leq n} (1 + X_p^{-1} X_q^{-1} - X_p^{-1}) \frac{1}{(1-X_1)(1-X_2) \dots (1-X_n)} \det_{1 \leq i, j \leq n} \left(\frac{X_i}{1-X_i} \right)^{j-1} \\ &= \frac{1}{(1-X_1)(1-X_2) \dots (1-X_n)} \prod_{1 \leq i < j \leq n} (1 + X_i^{-1} X_j^{-1} - X_i^{-1}) \left(\frac{X_j}{(1-X_j)} - \frac{X_i}{(1-X_j)} \right) \\ &= \frac{1}{(1-X_1)(1-X_2) \dots (1-X_n)} \prod_{1 \leq i < j \leq n} \frac{(X_j - X_i)(1 - X_j + X_i X_j)}{X_i(1-X_i)X_j(1-X_j)} \\ &= \prod_{i=1}^n \frac{1}{X_i^{n-1}(1-X_i)^n} \prod_{1 \leq i < j \leq n} (X_j - X_i)(1 - X_j + X_i X_j), \end{aligned}$$

where the Vandermonde determinant evaluation is used again. We summerize the result in the following theorem.

Theorem 3. *The coefficient of $X_1^{k_1} X_2^{k_2} \dots X_n^{k_n}$ in*

$$\prod_{i=1}^n \frac{1}{X_i^{n-1}(1-X_i)^n} \prod_{1 \leq i < j \leq n} (X_j - X_i)(1 - X_j + X_i X_j) \quad (6.4)$$

is equal to the number of monotone triangles with bottom row $(k_1, \dots, k_n)$ if $0 \leq k_1 < k_2 < \dots < k_n$ and $k_i \in \mathbb{Z}$, where (6.4) is interpreted as a formal laurent series, $\frac{1}{1-X_i} = \sum_{j=0}^{\infty} X_i^j$.

Consequently, the enumeration of $n \times n$ alternating sign matrices amounts to compute the constant term of

$$\prod_{i=1}^n \frac{1}{X_i^{n+i-2}(1-X_i)^n} \prod_{1 \leq i < j \leq n} (X_j - X_i)(1 - X_j + X_i X_j).$$

This is because monotone triangles with bottom row $(0, 1, \dots, n-1)$ correspond to $n \times n$ alternating sign matrices. Zeilberger [11] has used constant term identities to give the first proof of the alternating sign matrices theorem. (His identities are different from our result.)

Note that, if, for instant, we choose $n = 3$ in the generating function in Theorem 3 and consider the coefficient of $X_1^3 X_2^2 X_3$ we obtain -1 , which is obviously not the number of monotone triangles with bottom row $(3, 2, 1)$, since there exists no monotone triangle with this property. This coefficient is of course the values of $\alpha(3; 3, 2, 1)$. On the other hand, if we consider monomials $X_1^{k_1} \dots X_n^{k_n}$ with negative exponents then their

coefficients are not equal to $\alpha(n; k_1, \dots, k_n)$: for example the coefficient of $X_1^{-1}X_2^2X_3^3$ is 7 and this is not $\alpha(3; -1, 2, 3) = 23$. (In order to compute the number of monotone triangles with bottom row $(-1, 2, 3)$ using the generating function from Theorem 3, one can make use of the fact that $\alpha(3; -1, 2, 3) = \alpha(3; -1+c, 2+c, 3+c)$ for all integers c .)

Next we derive an analog generating function for halved monotone triangles with prescribed bottom row. Observe that Theorem 1 is equivalent to

$$\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil}) = \left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (E_{k_p}^{-1} + E_{k_q}^{-1} - \text{id})(E_{k_p}^{-1}E_{k_q}^{-1} + \text{id} - E_{k_q}^{-1}) \right) \bar{\gamma}(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

where

$$\bar{\gamma}(n, x; k_1, \dots, k_{(n+1)/2}) = \prod_{1 \leq i < j \leq (n+1)/2} \frac{(k_j - k_i)(2x + 2 - n - k_i - k_j)}{(j - i)(j + i - 1)}$$

if n is odd and

$$\bar{\gamma}(n, x; k_1, \dots, k_{n/2}) = \prod_{1 \leq i < j \leq n/2} \frac{(k_j - k_i)(2x + 4 - n - k_i - k_j)}{(j - i)(j + i)} \prod_{i=1}^{n/2} \frac{x + 2 - n/2 - k_i}{i}$$

if n is even. Here, we define $\bar{\gamma}_c(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ to be equal to $\bar{\gamma}(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$ if $k_l \leq c$ for all l and zero elsewhere. Then

$$\gamma(n, x; k_1, \dots, k_{\lceil n/2 \rceil}) = \left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (E_{k_p}^{-1} + E_{k_q}^{-1} - \text{id})(E_{k_p}^{-1}E_{k_q}^{-1} + \text{id} - E_{k_q}^{-1}) \right) \bar{\gamma}_c(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$$

for all $(k_1, \dots, k_{\lceil n/2 \rceil})$ with $k_l \leq c$. We compute the generating function

$$\sum_{(k_1, \dots, k_{\lceil n/2 \rceil}) \leq (n-1+c, \dots, n-1+c)} X_1^{k_1} \dots X_{\lceil n/2 \rceil}^{k_{\lceil n/2 \rceil}} \times \left(\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (E_{k_p}^{-1} + E_{k_q}^{-1} - \text{id})(E_{k_p}^{-1}E_{k_q}^{-1} + \text{id} - E_{k_q}^{-1}) \right) \bar{\gamma}_c(n, x; k_1, \dots, k_{\lceil n/2 \rceil}). \quad (6.5)$$

Let

$$\prod_{1 \leq p < q \leq \lceil n/2 \rceil} (Y_p^{-1} + Y_q^{-1} - 1)(Y_p^{-1}Y_q^{-1} + 1 - Y_q^{-1}) = \sum_{\substack{(j_1, \dots, j_{\lceil n/2 \rceil}) \\ -n+1 \leq j_i \leq 0}} b(j_1, \dots, j_{\lceil n/2 \rceil}) Y_1^{j_1} \dots Y_{\lceil n/2 \rceil}^{j_{\lceil n/2 \rceil}}.$$

Thus, the generating function (6.5) is equal to

$$\sum_{(k_1, \dots, k_{\lceil n/2 \rceil}) \leq (n-1+c, \dots, n-1+c)} \sum_{\substack{(j_1, \dots, j_{\lceil n/2 \rceil}) \\ -n+1 \leq j_i \leq 0}} b(j_1, \dots, j_{\lceil n/2 \rceil}) X_1^{k_1} \dots X_{\lceil n/2 \rceil}^{k_{\lceil n/2 \rceil}} \\ \times \overline{\gamma}_c(n, x; k_1 + j_1, \dots, k_{\lceil n/2 \rceil} + j_{\lceil n/2 \rceil}). \quad (6.6)$$

Again we set $(l_1, \dots, l_{\lceil n/2 \rceil}) = (k_1 + j_1, \dots, k_{\lceil n/2 \rceil} + j_{\lceil n/2 \rceil})$. Consequently, (6.6) is equal to

$$\sum_{\substack{(j_1, \dots, j_{\lceil n/2 \rceil}) \\ -n+1 \leq j_i \leq 0}} b(j_1, \dots, j_{\lceil n/2 \rceil}) X_1^{-j_1} \dots X_{\lceil n/2 \rceil}^{-j_{\lceil n/2 \rceil}} \\ \times \sum_{(l_1, \dots, l_{\lceil n/2 \rceil}) \leq (n-1+j_1+c, \dots, n-1+j_{\lceil n/2 \rceil}+c)} \overline{\gamma}_c(n, x; l_1, \dots, l_{\lceil n/2 \rceil}) X_1^{l_1} \dots X_{\lceil n/2 \rceil}^{l_{\lceil n/2 \rceil}}. \quad (6.7)$$

Since $j_l \geq -n+1$ for all l and $\overline{\gamma}_c(n, x; l_1, \dots, l_{\lceil n/2 \rceil}) = 0$ if $l_i > c$ for an i , (6.7) is equal to

$$\sum_{\substack{(j_1, \dots, j_{\lceil n/2 \rceil}) \\ -n+1 \leq j_i \leq 0}} b(j_1, \dots, j_{\lceil n/2 \rceil}) X_1^{-j_1} \dots X_{\lceil n/2 \rceil}^{-j_{\lceil n/2 \rceil}} \\ \times \sum_{(l_1, \dots, l_{\lceil n/2 \rceil}) \leq (c, \dots, c)} \overline{\gamma}_c(n, x; l_1, \dots, l_{\lceil n/2 \rceil}) X_1^{l_1} \dots X_{\lceil n/2 \rceil}^{l_{\lceil n/2 \rceil}} \\ = \prod_{1 \leq p < q \leq \lceil n/2 \rceil} (X_p + X_q - 1)(X_p X_q + 1 - X_q) \\ \times \sum_{(l_1, \dots, l_{\lceil n/2 \rceil}) \leq (c, \dots, c)} \overline{\gamma}_c(n, x; l_1, \dots, l_{\lceil n/2 \rceil}) X_1^{l_1} \dots X_{\lceil n/2 \rceil}^{l_{\lceil n/2 \rceil}}. \quad (6.8)$$

In the last expression $\overline{\gamma}_c$ can be replaced by $\overline{\gamma}$. The following lemma provides us with determinantal expressions for $\overline{\gamma}(n, x; k_1, \dots, k_{\lceil n/2 \rceil})$.

Lemma 13. (1)

$$\det_{1 \leq i, j \leq n} \binom{k_i + j - 1}{2j - 1} = \prod_{1 \leq i < j \leq n} \frac{(k_j - k_i)(k_i + k_j)}{(j - i)(j + i)} \prod_{i=1}^n \frac{k_i}{i}$$

(2)

$$\det_{1 \leq i, j \leq n} \binom{k_i + j - 3/2}{2j - 2} = \prod_{1 \leq i < j \leq n} \frac{(k_j - k_i)(k_i + k_j)}{(j - i)(j + i - 1)}$$

Proof. We only prove (1) since the proof of (2) is similar. First observe that

$$\binom{k_i + j - 1}{2j - 1} = \frac{k_i}{(2j - 1)!} \prod_{l=1}^{j-1} (k_i^2 - l^2).$$

Thus, the determinant in (1) is equal to

$$\prod_{i=1}^{n-1} \frac{1}{(2i+1)!} \prod_{i=1}^n k_i \det_{1 \leq i, j \leq n} \left(\prod_{l=1}^{j-1} (k_i^2 - l^2) \right).$$

The assertion follows from

$$\det_{1 \leq i, j \leq n} p_j(Y_i) = \prod_{1 \leq i < j \leq n} (Y_j - Y_i),$$

where $p_j(Y)$ is a polynomial in Y of degree $j-1$ whose leading coefficient is 1. This is a consequence of the Vandermonde determinant evaluation. $\square$

Lemma 13 implies that

$$\overline{\gamma}(n, x; k_1, \dots, k_{(n+1)/2}) = (-1)^{\binom{(n+1)/2}{2}} \det_{1 \leq i, j \leq (n+1)/2} \binom{k_i + j + n/2 - x - 5/2}{2j-2} \quad (6.9)$$

if n is odd and

$$\overline{\gamma}(n, x; k_1, \dots, k_{n/2}) = (-1)^{\binom{(n+2)/2}{2}} \det_{1 \leq i, j \leq n/2} \binom{k_i + j + n/2 - x - 3}{2j-1} \quad (6.10)$$

if n is even. If we use these determinantal presentations for $\overline{\gamma}$ in (6.8), we obtain the following generating function

$$\begin{aligned} & \prod_{1 \leq p < q \leq (n+1)/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \\ & \times \det_{1 \leq i, j \leq (n+1)/2} \left(\sum_{l_i = -\infty}^c \binom{l_i + j + n/2 - x - 5/2}{2j-2} X_i^{l_i} \right) \end{aligned} \quad (6.11)$$

if n is odd, and

$$\begin{aligned} & \prod_{1 \leq p < q \leq n/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \\ & \times \det_{1 \leq i, j \leq n/2} \left(- \sum_{l_i = -\infty}^c \binom{l_i + j + n/2 - x - 3}{2j-1} X_i^{l_i} \right) \end{aligned} \quad (6.12)$$

if n is even.

If we choose $c = x + 1/2 - n/2$ in case that n is odd and $c = x + 2 - n/2$ in case that n is even, the determinants in the expression above simplify. This follows from the following identities.

$$\begin{aligned} & \sum_{l=-\infty}^c \binom{l + j - c + z}{2j-2} X^l = \frac{X^{j+c-z-2}}{(2j-2)!} \frac{d}{dX^{2j-2}} \left(\sum_{l=-\infty}^c X^{l+j-c+z} \right) \\ & = \frac{X^{j+c-z-2}}{(2j-2)!} \frac{d}{dX^{2j-2}} \left(\frac{X^{j+z+1}}{X-1} \right) = \frac{X^{j+c-z-2}}{(2j-2)!} \frac{d}{dX^{2j-2}} \left(\frac{1}{X-1} \right) = \frac{X^{j+c-z-2}}{(X-1)^{2j-1}} \end{aligned} \quad (6.13)$$

where $z = -j - 1, -j, \dots, j - 3$. (Note that the identity is true for all j if $z = -2$.) Similarly,

$$\sum_{l=-\infty}^c \binom{l+j-c+z}{2j-1} X^l = -\frac{X^{j+c-z-1}}{(X-1)^{2j}} \quad (6.14)$$

where $z = -j - 1, -j, \dots, j - 2$. (This identity is true for all j if $z = -2, -1$.)

We first consider the case that n is odd. By (6.13) ($z = -2$) the generating function (6.11) is equal to

$$\begin{aligned} & \prod_{1 \leq p < q \leq (n+1)/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \det_{1 \leq i, j \leq (n+1)/2} \left(\frac{X_i^{j+c}}{(X_i - 1)^{2j-1}} \right) \\ &= \prod_{1 \leq p < q \leq (n+1)/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \prod_{i=1}^{(n+1)/2} \frac{X_i^{c+1}}{(X_i - 1)} \det_{1 \leq i, j \leq (n+1)/2} \left(\frac{X_i}{(X_i - 1)^2} \right)^{j-1}. \end{aligned}$$

The Vandermonde determinant evaluation shows that this is equal to

$$\begin{aligned} & \prod_{1 \leq p < q \leq (n+1)/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \prod_{i=1}^{(n+1)/2} \frac{X_i^{c+1}}{(X_i - 1)} \\ & \quad \times \prod_{1 \leq i < j \leq (n+1)/2} \left(\frac{X_j}{(X_j - 1)^2} - \frac{X_i}{(X_i - 1)^2} \right) \\ &= \prod_{1 \leq i < j \leq (n+1)/2} (X_j - X_i)(X_i + X_j - 1)(X_i X_j - 1)(1 - X_j + X_i X_j) \prod_{i=1}^{(n+1)/2} \frac{X_i^{x+3/2-n/2}}{(X_i - 1)^n}. \end{aligned} \quad (6.15)$$

Finally we consider the case that n is even. By (6.14) ($z = -1$) the generating function (6.12) is equal to

$$\begin{aligned} & \prod_{1 \leq p < q \leq n/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \det_{1 \leq i, j \leq n/2} \left(\frac{X_i^{j+c}}{(X_i - 1)^{2j}} \right) = \\ & \prod_{1 \leq p < q \leq n/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \prod_{i=1}^{n/2} \frac{X_i^{c+1}}{(X_i - 1)^2} \det_{1 \leq i, j \leq n/2} \left(\frac{X_i}{(X_i - 1)^2} \right)^{j-1} \end{aligned}$$

The Vandermonde determinant evaluation now shows that this is equal to

$$\begin{aligned}
& \prod_{1 \leq p < q \leq n/2} (1 - X_p - X_q)(X_p X_q + 1 - X_q) \prod_{i=1}^{n/2} \frac{X_i^{c+1}}{(X_i - 1)^2} \\
& \quad \times \prod_{1 \leq i < j \leq n/2} \left(\frac{X_j}{(X_j - 1)^2} - \frac{X_i}{(X_i - 1)^2} \right) \\
& = \prod_{1 \leq i < j \leq n/2} (X_j - X_i)(X_i + X_j - 1)(X_i X_j - 1)(1 - X_j + X_i X_j) \prod_{i=1}^{n/2} \frac{X_i^{x+3-n/2}}{(X_i - 1)^n}. \quad (6.16)
\end{aligned}$$

In this case the generating functions in (6.15) and (6.16) are understood as formal laurent series in $1/X_i$, i.e. $1/(X_i - 1) = \sum_{j=-\infty}^{-1} X_i^j$.

REFERENCES

- [1] D. M. Bressoud, *Proof and confirmations, The story of the alternating sign matrix conjecture*, Cambridge University Press, Cambridge, 1999.
- [2] I. Fischer, The number of monotone triangles with prescribed bottom row, *Adv. in Appl. Math.* **37** (2006), no. 2, 249 – 267.
- [3] I. Fischer, A new proof of the refined alternating sign matrix theorem, *J. Combin. Theory Ser. A* **114** (2007), no. 2, 253 – 264.
- [4] G. Kuperberg, Another proof of the alternating-sign matrix conjecture, *Internat. Math. Res. Notices.* **1996**, no. 3, 139 – 150.
- [5] G. Kuperberg, Symmetry classes of alternating-sign matrices under one roof, *Ann. of Math. (2)* **156** (2002), no. 3, 835 – 866.
- [6] W. H. Mills, D. P. Robbins and H. Rumsey, Proof of the Macdonald conjecture, *Invent. Math.* **66** (1982), no. 1, 73 – 87.
- [7] W. H. Mills, D. P. Robbins and H. Rumsey, Alternating sign matrices and descending plane partitions, *J. Combin. Theory Ser. A* **34** (1983), no. 3, 340 – 359.
- [8] R. A. Proctor, Odd symplectic groups, *Invent. math.* **92** (1988), 307 – 332.
- [9] A. V. Razumov and Yu. G. Stroganov, On refined enumerations of some symmetry classes of ASMs, *Theor.Math.Phys.* **141** (2004) 1609-1630; *Teor.Mat.Fiz.* **141** (2004) 323-347
- [10] D. P. Robbins, Symmetry classes of alternating sign matrices, preprint, [math.CO/0008045](#)
- [11] D. Zeilberger, Proof of the alternating sign matrix conjecture, The Foata Festschrift, *Electron. J. Combin.* **3** (1996), no. 2, R13, 84 pp
- [12] D. Zeilberger, Proof of the refined alternating sign matrix conjecture, *New York J. Math.* **2** (1996), 59 – 68.